

Astroparticle Physics

2020/21

Tuesday 13:30 - 15:15

Thursday 13:30 - 15:15

- lectures
- student presentations
- oral exam, ca. 45 min

Jörg R. Hörandel


HG 02.728

j.horandel@astro.ru.nl

<http://particle.astro.ru.nl/goto.html?astropart2021>




birth of cosmic rays
CRs: supernova remnants
neutrinos: e.g. Sun (lecture 8)



propagation of CRs in the Heliosphere (lecture 9)




propagation of CRs in the Galaxy
interactions with ISM (lecture 9)



The Earth's Magnetic Field

propagation of CRs in the Earth magnetic field (lecture 9)



CRs in the atmosphere (lecture 10)


CRs at the top of the atmosphere (lecture 10)



CRs underground (lecture 11)
neutrino oscillations (lecture 11+12)

Particles and the Cosmos


$\sim 1000/m^2 \text{ s}$



energy spectrum
of cosmic rays
extends to
extremely high
energies:
 $10^{20} \text{ eV} \sim 16 \text{ J}$

Origin of cosmic rays

multi messenger technique



RESEARCH ARTICLE SUMMARY

NEUTRINO ASTROPHYSICS


Multimessenger observations of a flaring blazar coincident with high-energy neutrino IceCube-170922A

The IceCube Collaboration, *Fermi*-LAT, MAGIC, *AGILE*, ASAS-SN, HAWC, H.E.S.S., *INTEGRAL*, Kanata, Kiso, Kapteyn, Liverpool Telescope, Subaru, *Swift*/*NuSTAR*, VERITAS, and VLA/17B-403 teams*†

INTRODUCTION: Neutrinos are tracers of cosmic-ray acceleration: electrically neutral and traveling at nearly the speed of light, they can escape the densest environments and may be traced back to their source of origin. High-energy neutrinos are expected to be produced in blazars: intense extragalactic radio, optical, x-ray, and, in some cases, γ -ray sources characterized by relativistic jets of plasma pointing close to our line of sight. Blazars are among the most powerful objects in the Universe and are widely speculated to be sources of high-energy cosmic rays. These cosmic rays generate high-energy neutrinos and γ -rays, which are produced when the cosmic rays accelerated in the jet interact with nearby gas or photons. On 22 September 2017, the cubic-kilometer IceCube Neutrino Observatory detected a ~ 290 -TeV neutrino from a direction consistent with the flaring γ -ray blazar TXS 0506+056. We report the details of this observation and the results of a multiwavelength follow-up campaign.

RATIONALE: Multimessenger astronomy aims for globally coordinated observations of cosmic rays, neutrinos, gravitational waves, and electromagnetic radiation across a broad range of wavelengths. The combination is expected to yield crucial information on the mechanisms

mic rays. The discovery of an extraterrestrial diffuse flux of high-energy neutrinos, announced by IceCube in 2013, has characteristic properties that hint at contributions from extragalactic sources, although the individual sources remain as yet unidentified. Continuously monitoring the entire sky for astrophysical neu-



Multimessenger observations of blazar TXS 0506+056. The 50% and 90% containment contours for the neutrino IceCube

trinos, IceCube provides real-time triggers for observatories around the world measuring γ -rays, x-rays, optical, radio, and gravitational waves, allowing for the potential identification of even rapidly fading sources.

RESULTS: A high-energy neutrino-induced muon track was detected on 22 September 2017, automatically generating an alert that was

ON OUR WEBSITE

Read the full article at <http://dx.doi.org/10.1126/science.aat1378>

distributed worldwide within 1 min of detection and prompted follow-up searches by telescopes over a broad range of wavelengths. On 28 September 2017, the *Fermi* Large Area


Telescope Collaboration reported that the direction of the neutrino was coincident with a cataloged γ -ray source, 0.1° from the neutrino direction. The source, a blazar known as TXS 0506+056 at a measured redshift of 0.34, was in a flaring state at the time with enhanced γ -ray activity in the GeV range. Follow-up observations by imaging atmospheric Cherenkov telescopes, notably the Major Atmospheric Gamma Imaging Cherenkov (MAGIC) telescopes, revealed periods where the detected γ -ray flux from the blazar reached energies up to 400 GeV. Measurements of the source have also been completed at x-ray, optical, and radio wavelengths. We have investigated models associating neutrino and γ -ray production and find that correlation of the neutrino with the flare of TXS 0506+056 is statistically significant at the level of 3 standard deviations (σ). On the basis of the redshift of TXS 0506+056, we derive constraints for the muon-neutrino luminosity for this source and find them to be similar to the luminosity observed in γ -rays.

CONCLUSION: The energies of the γ -rays and the neutrino indicate that blazar jets may accelerate cosmic rays to at least several PeV. The observed association of a high-energy neutrino with a blazar during a period of enhanced γ -ray emission suggests that blazars may indeed be one of the long-sought sources of cosmic high-energy


Downloaded from <http://science.sciencemag.org/> on December 11, 2019

DATA FOR THE R-BAND IN MAGNIFIED VIEW



Follow-up of GW170817 with PAO (neutrinos)



LIGO



Follow-up of GW170817 with PAO (neutrinos)



THE ASTROPHYSICAL JOURNAL LETTERS, 848:L12 (59pp), 2017 October 20

<https://doi.org/10.3847/2041-8213/aa91c9>

© 2017, The American Astronomical Society. All rights reserved.

OPEN ACCESS



Multi-messenger Observations of a Binary Neutron Star Merger

LIGO Scientific Collaboration and Virgo Collaboration, Fermi GBM, INTEGRAL, IceCube Collaboration, AstroSat Cadmium Zinc Telluride Imager Team, IPN Collaboration, The Insight-Hxmt Collaboration, ANTARES Collaboration, The Swift Collaboration, AGILE Team, The 1M2H Team, The Dark Energy Camera GW-EM Collaboration and the DES Collaboration, The DLT40 Collaboration, GRAWITA: GRAvitational Wave Inaf TeAm, The Fermi Large Area Telescope Collaboration, ATCA: Australia Telescope Compact Array, ASKAP: Australian SKA Pathfinder, Las Cumbres Observatory Group, OzGrav, DWF (Deeper, Wider, Faster Program), AST3, and CAASTRO Collaborations, The VINROUGE Collaboration, MASTER Collaboration, J-GEM, GROWTH, JAGWAR, CaltechNRAO, TTU-NRAO, and NuSTAR Collaborations, Pan-STARRS, The MAXI Team, TZAC Consortium, KU Collaboration, Nordic Optical Telescope, ePESSTO, GROND, Texas Tech University, SALT Group, TOROS: Transient Robotic Observatory of the South Collaboration, The BOOTES Collaboration, MWA: Murchison Widefield Array, The CALET Collaboration, IKI-GW Follow-up Collaboration, H.E.S.S. Collaboration, LOFAR Collaboration, LWA: Long Wavelength Array, HAWC Collaboration, **The Pierre Auger Collaboration**, ALMA Collaboration, Euro VLBI Team, Pi of the Sky Collaboration, The Chandra Team at McGill University, DFN: Desert Fireball Network, ATLAS, High Time Resolution Universe Survey, RIMAS and RATIR, and SKA South Africa/MeerKAT (See the end matter for the full list of authors.)

Received 2017 October 3; revised 2017 October 6; accepted 2017 October 6; published 2017 October 16

Follow-up of GW170817 with PAO (neutrinos)

THE ASTROPHYSICAL JOURNAL LETTERS, 850:L35 (18pp), 2017 December 1
© 2017. The American Astronomical Society.

<https://doi.org/10.3847/2041-8213/aa9aed>




OPEN ACCESS

Search for High-energy Neutrinos from Binary Neutron Star Merger GW170817 with ANTARES, IceCube, and the Pierre Auger Observatory

ANTARES Collaboration, IceCube Collaboration, The Pierre Auger Collaboration, and LIGO Scientific Collaboration and Virgo Collaboration

PAO in pre-defined +/- 500 s window as sensitive as IceCube

GW170817 Neutrino limits (fluence per flavor: $\nu_x + \bar{\nu}_x$)



The γ -ray spectrum of the core of Centaurus A as observed with H.E.S.S. and *Fermi*-LAT

ABSTRACT

Centaurus A (Cen A) is the nearest radio galaxy discovered as a very-high-energy (VHE; 100 GeV–100 TeV) γ -ray source by the High Energy Stereoscopic System (H.E.S.S.). It is a faint VHE γ -ray emitter, though its VHE flux exceeds both the extrapolation from early *Fermi*-LAT observations as well as expectations from a (misaligned) single-zone synchrotron-self Compton (SSC) description. The latter satisfactorily reproduces the emission from Cen A at lower energies up to a few GeV. New observations with H.E.S.S., comparable in exposure time to those previously reported, were performed and eight years of *Fermi*-LAT data were accumulated to clarify the spectral characteristics of the γ -ray emission from the core of Cen A. The results allow us for the first time to achieve the goal of constructing a representative, contemporaneous γ -ray core spectrum of Cen A over almost five orders of magnitude in energy. Advanced analysis methods, including the template fitting method, allow detection in the VHE range of the core with a statistical significance of 12σ on the basis of 213 hours of total exposure time. The spectrum in the energy range of 250 GeV–6 TeV is compatible with a power-law function with a photon index $\Gamma = 2.52 \pm 0.13_{\text{stat}} \pm 0.20_{\text{sys}}$. An updated *Fermi*-LAT analysis provides evidence for spectral hardening by $\Delta\Gamma \approx 0.4 \pm 0.1$ at γ -ray energies above $2.8^{+1.0}_{-0.6}$ GeV at a level of 4.0σ . The fact that the spectrum hardens at GeV energies and extends into the VHE regime disfavour a single-zone SSC interpretation for the overall spectral energy distribution (SED) of the core and is suggestive of a new γ -ray emitting component connecting the high-energy emission above the break energy to the one observed at VHE energies. The absence of significant variability at both GeV and TeV energies does not yet allow disentanglement of the physical nature of this component, though a jet-related origin is possible and a simple two-zone SED model fit is provided to this end.

Key words. gamma rays: galaxies – radiation mechanisms: non-thermal





Fig. 3. SED of Cen A core with model fits as described in text. The red curve corresponds to an SSC component designed to fit the radio to sub-GeV data. The blue curve corresponds to a second SSC component added to account for the highest energy data. The black curve corresponds to the sum of the two components. SED points as derived from H.E.S.S. and *Fermi*-LAT data in this paper are shown with open circles. Observations from the radio band to the MeV γ -ray band are from TANAMI (\diamond), SEST (\blacktriangle), JCMT (\triangleright), MIDI (∇), NAOS/CONICA (\triangleleft), NICMOS (\square), WFPC2 (\blacklozenge), *Suzaku* (\triangle), OSSE/COMPTEL (\blacksquare). The acronyms are described in Appendix B.



Observation of a large-scale anisotropy in the arrival directions of cosmic rays above 8×10^{18} eV

The Pierre Auger Collaboration*†



Anisotropy detected at >5.2 sigma dipole amplitude 6.5%





Fig. 3. Map showing the fluxes of particles in galactic coordinates. Sky map in galactic coordinates showing the cosmic-ray flux for $E \geq 8$ EeV smoothed with a 45° top-hat function. The galactic center is at the origin. The cross indicates the measured dipole direction; the contours denote the 68% and 95% confidence level regions. The dipole in the 2MRS galaxy distribution is indicated. Arrows show the deflections expected for a particular model of the galactic magnetic field (8) on particles with $E/Z = 5$ or 2 EeV.

Astroparticle Physics

messengers from the Universe



gravitational waves

(charged) cosmic rays

gamma rays

3-K microwave background

neutrinos

dark matter

GZK?

γ_{3K}


Literature

Particles & Cosmos: Stanev

Astroparticle Physics:

Tom Gaisser, Cosmic rays and particle physics
Cambridge University Press (2016)

+ primary literature (journal articles)



Astroparticle Physics

2020/21

1. Historical introduction - basic properties of cosmic rays
2. Hadronic interactions and accelerator data
3. Cascade equations
4. Electromagnetic cascades
5. Extensive air showers
6. Detectors for extensive air showers
7. High-energy cosmic rays and the knee in the energy spectrum of cosmic rays
8. Radio detection of extensive air showers
9. Acceleration, Astrophysical accelerators and beam dumps
10. Extragalactic propagation of cosmic rays
11. Ultra-high-energy energy cosmic rays
12. Astrophysical gamma rays and neutrinos
13. Neutrino astronomy
14. Gamma-ray astronomy

Student talks

- **Students will present selected topics, based on journal publications.**
- **Learn how to derive information from primary literature.**
- **Presentation followed by discussion and questions.**
- **60 min presentation, 15 min discussion**
- **You are expected to participate in discussions and ask questions.**
- **Your presentation + interaction will be part of your grade.**

Student talks

- **Air showers - Matthews Heitler model**
- **Radio detection of air showers**
- **CR anisotropy at TeV energies, IceCube/Top, HAWC**
- **the knee in the energy spectrum of cosmic rays**
- **Detectors for UHE CRs, Auger, TA**
- **Auger proton-air cross section**
- **GZK effect and the end of the CR spectrum, Auger, TA**
- **CR mass composition at highest energies, Auger, TA**
- **CR anisotropy at highest energies, Auger, TA**
- **IceCube neutrino astronomy**
- **KM3NeT project ARCA+ORCA**
- **H.E.S.S. TeV gamma-ray astronomy galactic center emission**
- **H.E.S.S. TeV gamma-ray astronomy galactic plane survey**
- **Cherenkov Telescope Array - CTA**
- **XENON dark matter search**
- **LIGO + Virgo gravitational waves**

break until 14:30


lecture 1

Historical introduction
Basic properties of Cosmic Rays

Discovery of Radioactivity




Henri Becquerel



Marie & Pierre Curie

Nobel Prize
1903




Ein neues Elektrometer für statische Ladungen.

Dritte Mitteilung¹⁾.

Von Th. Wulf.

Mitteilung enthält einige
weiter beschriebenen Appa-
raturerhöhung seiner Transport-

a new electrometer
for static charges





**Sir J.J.Thomson
Nobel Prize 1906**




Conduction of electricity through gases (1928):

It would be one of the romances of science if these obscure and prosaic minute leakages of electricity from well-insulated bodies should be the means by which the most fundamental problems in the evolution of the cosmos came to be investigated.

Credit Alex MacDonald

Detector used by Wilson to investigate ionization of air



“the continuous production of ions in dust-free air could be explained as being due to radiation from sources outside our atmosphere, possibly radiation like Röntgen rays or cathode rays, but of enormously greater penetrating power”

C T R Wilson, Proc Roy Soc A 68 (1901) 151

on the origin of gamma radiation in the atmosphere

Über den Ursprung der in der Atmosphäre vorhandenen γ -Strahlung.

Von Th. Wulf.

Tabelle I.

Strahlung der Wände von Gebäuden.

| Ort | Material | Alter | Strahlung Ionen pro cm ² u. Sekunde |
|---|-----------------|-----------|--|
| Abtei Maria Laach bei Andernach a. Rh. | Vulkanisch Tuff | 50 Jahre | 13,7 |
| Valkenburg, Colleg, Holland-L. | | | |
| Löwen, Colleg, Belgien | Ziegelsteine | 15 " | 3,7 |
| Namur, Colleg N. D. de la paix, Belgien | Ziegelsteine | — | 8,0 |
| Wynandsrade Kasteel, Holland | Ziegelsteine | ca. 100 | 3,7 |
| | Ziegelsteine | 200 Jahre | 0,0 |

Man kann den Inhalt dieser Arbeit kurz so zusammenfassen. Es wird über Versuche berichtet, welche beweisen, daß an dem Beobachtungsort die durchdringende Strahlung von primär radioaktiven Substanzen verursacht wird, welche in den obersten Erdschichten liegen, bis etwa 1 m unter der Oberfläche.

Wenn ein Teil der Strahlung aus der Atmosphäre stammt, so ist er doch so klein, daß er sich mit den gebrauchten Mitteln nicht nachweisen ließ.

Die zeitlichen Schwankungen in der γ -Strahlung

leben
de
ie
N

Nur in dem alten holländischen Kasteel Wynandsrade, vor fast 200 Jahren aus Ziegelsteinen erbaut, zeigte sich kein Unterschied in der Strahlung im Zimmer und im Freien. — Am stärksten war die Strahlung in Maria Laach in einem

the radiation originates from the soil maybe a small contribution from the atmosphere



~1910



Theodor Wulf

**1909: Soddy & Russel:
attenuation of gamma rays
follows an exponential law**

$$I = I_0 e^{-\mu L}$$

Discovery of Cosmic Rays

Viktor Franz Hess

7. August 1912

Early cosmic-ray work published in German

Jörg R. Hörandel

Citation: *AIP Conf. Proc.* **1516**, 52 (2013); doi: 10.1063/1.4792540




FIGURE 1. *Left:* Electrometer after Th. Wulf [5]. *Right:* Two grandsons of V.F. Hess revealing a plaque to commemorate the discovery of cosmic rays on August 7th, 2012, close to the presumed landing site of V.F. Hess in Pieskow close to Berlin. It reads: "To commemorate the discovery of cosmic rays. On 7 August 1912 landed the Austrian physicist Victor F. Hess with a hydrogen balloon close to Pieskow. On the journey from Lower-Bohemia he reached an altitude of 5300 m and he proved the existence of a penetrating, ionizing radiation from outer space. For the discovery of cosmic rays V.F. Hess has been awarded the Nobel Prize in Physics in 1936. The participants of the symposium '100 years cosmic rays', Bad Saarow-Pieskow, 7 August 2012".

Aeronautisches Gelände im Wiener Prater, wo seine ersten Freiballon-Forschungsfahrten u schichtfliehe Mu

Hess on

Route des Entdeckungsfluges der kosmischen Strahlung.

Aus der Abteilung für Geophysik, Meteorologie
und Erdmagnetismus:

Viktor F. Hess (Wien), Über Beobachtungen
der durchdringenden Strahlung bei sieben
Freiballonfahrten.

on the observation of
the penetrating
radiation during 7
balloon campaigns

Im Vorjahre habe ich bereits Gelegenheit
gehabt, zwei Ballonfahrten zur Erforschung
der durchdringenden Strahlung zu unterneh-
men: über die erste Fahrt

7. Fahrt (7. August 1912).

Ballon: „Böhmen“ (1680 cbm Wasserstoff).
Meteorolog. Beobachter: E. Wolf.

Führer: Hauptmann W. Hoffory.
Luftelektr. Beobachter: V. F. Hess.

| | | Mittlere Höhe | | Beobachtete Strahlung | | | | Temp. | Relat. Feucht. Proz. |
|----|---------------|---------------|--------------|-----------------------|-------------|-------------|--------------------|---|----------------------------|
| | | absolut | relativ m | Apparat 1 | Apparat 2 | Apparat 3 | | | |
| | | | | φ_1 | φ_2 | φ_3 | reduz. φ_3 | | |
| 1 | 15h 15—16h 15 | 156 | 0 | 17,3 | 12,9 | — | — | 1½ Tag vor dem Auf- stiege (in Wien) | |
| 2 | 16h 15—17h 15 | 156 | 0 | 15,9 | 11,0 | 18,4 | 18,4 | | |
| 3 | 17h 15—18h 15 | 156 | 0 | 15,8 | 11,2 | 17,5 | 17,5 | | |
| 4 | | 1700 | 1400 | 15,8 | 14,4 | — | — | | |
| | | 2750 | 2500 | 17,3 | 17,3 | — | — | | |
| | | 3850 | 3600 | 19,8 | 19,8 | — | — | | |
| 7 | | 4800 | 4700 | 40,7 | 36,7 | — | — | | |
| | | (4400) | (3300) | — | — | — | — | | |
| 8 | 10h 45—11h 15 | 4400 | 4200 | 28,1 | 22,7 | — | — | | |
| 9 | 11h 15—11h 45 | 1300 | 1200 | (9,7) | 11,5 | — | — | | |
| 10 | 11h 45—12h 10 | 250 | 150 | 11,9 | 10,7 | — | — | | |
| 11 | 12h 25—13h 12 | 140 | 0 | 15,0 | 11,6 | — | — | | |

hydrogen!

altitude

intensity

Aus der Abteilung für Geophysik, Meteorologie
und Erdmagnetismus:

Viktor F. Hess (Wien), Über Beobachtungen
der durchdringenden Strahlung bei sieben



V.F. Hess in 1936–37, on the occasion of Nobel prize.

Nobel Prize 1936

der Verringerung der radioaktiven Substanzen
der Atmosphäre zurückzuführen.

Die Ergebnisse der vorliegenden Beobachtungen scheinen am ehesten durch die Annahme erklärt werden zu können, daß eine Strahlung von sehr hoher Durchdringungskraft von oben her in unsere Atmosphäre eindringt, und auch noch in deren untersten Schichten einen Teil der in geschlossenen Gefäßen beobachteten Ionisation hervorruft. Die Intensität dieser Strahlung scheint zeitlichen Schwankungen unterworfen zu sein, welche bei einstündigen Ablesungsintervallen noch erkennbar sind. Da ich im Ballon weder bei Nacht noch bei einer Sonnenfinsternis eine Verringerung der Strahlung fand, so kann man wohl kaum die Sonne als Ursache dieser hypothetischen Strahlung ansehen, wenigstens solange man nur an eine direkte γ -Strahlung mit geradliniger Fortpflanzung denkt.

Daß die Zunahme der Strahlung erst jenseits 3000 m so stark merklich wird, ist nicht

erweitertes Beobachtungsmaterial wurde.



Neue Untersuchungen über die durchdringende Hesssche Strahlung.

Von E. Steinke in Königsberg i. Pr.

Absorption in the atmosphere

Intensity as function for different altitudes

intensity




atmospheric overburden

Über Schwankungen und Barometereffekt der kosmischen Ultrastrahlung im Meeresniveau.

Von E. Steinke in Königsberg i. Pr.

Barometric effect

Während d
stunden au
Dicke, teil
material e
teils perio
Schwanku
dische Sch
mäßige In
endlich pe
während c
kungen, in
für das Au
zeitlichen



with
ect

on of



Fig. 4. Zusammenhang zwischen Barometerstand und Ionisation (Jahresmittel der Stundenwerte; die Zahlen geben die Anzahl der Stundenwerte an).

si

Absorption in Lake Constance 1928




Fig. 5.





Fig. 6. Die „schwimmende“ Verankerung des Apparates.



Absorption in Lake Constance 1928

Ionization chamber with electrometer read-out
automatic each hour, up to 8 days




Fig. 1.
Ausbau des wasserdichten
Registrierapparates.






Fig. 1.

Absorption in Lake Constance 1928



equivalent depth (of water)
from top of the atmosphere

Fig. 6.



Hess

Steinke

Regener

**Three pioneers of Cosmic Ray research
Regener demonstrates his balloon electrometer
(Immenstaad/Lake Constance, August 1932).**

derung zu danken, die die ...
von An ...
8 ...

Kolhörster
A new electrometer

1) Oskar Taussig, ...
(The First World ...
vgl. auch „Elektrotechnik ...
des Elektrotechnischen ...


(3. Jahrgang 28. August 1925)

Ein neues Fadenelektrometer.

Von Werner Kolhörster.

Zu Messungen der durchdringenden Strahlung hatte ich für meine neuen Strahlungsapparate ein Fadenelektrometer konstruiert¹⁾, das ohne die bei derartigen Instrumenten notwendige Temperaturkompensation arbeitet. Da es sich auch für andere elektrostatische Messungen seiner Vorträge und allgemeinen Verwendbarkeit halber als geeignet erwies, so seien hier einige Angaben über die Instrumente²⁾ gemacht.

Prinzip: Als Gegenkraft gegen die elektrostatischen Abstößungskräfte dient allein die Biegeelastizität der feinen Quarzfäden, die die Form vertikal stehender, frei tragender Schlingen haben und deren Enden in einigen Millimetern Abstand voneinander an einem Metallblech befestigt sind, das in den Isolator eingesetzt wird. Entsprechend den Ein- und Zweifadenelektrometern kann man Systeme mit einer oder zwei kongruenten Schlingen verwenden, die von einem Mikroskop mit Okularmikrometer am Scheitel der Schlingen abgelesen werden. Lädt man das System, so tritt keine merkliche Formänderung der Schlingen ein, diese bewegen sich vielmehr in der Horizont-



ldung zu verändern. Eine feinere Änderung läßt sich durch Verschwenken der Schneiden erzielen, die um die längere Rechteckseite drehbar, mehr oder weniger den Fäden genähert



Kohlhörster - balloon flight 13. May 1934




Abb.12 Vor dem Aufstieg




Abb.19 Höhenballon "Bartsch von Sigsfeld"
Füllung am 13.5.34 rd. 4400 m³.

Kohlhörster - balloon flight 13. May 1934



Abb. 17

Dr. Schrenk




Abb. 1

Masuch

Measurements of the cosmic-ray intensity (Höhenstrahlung) up to 12000 m



Fig. 19. Regener recovering a balloon payload from a farm house.

3) Die Firma Gebr. Junghans, Schramberg, hat uns freundlicherweise diese schönen Zählwerke hergestellt.





Fig. 6. Registrierapparat mit Schutzgondel.

Das Wesen der Höhenstrahlung.

Von W. Bothe und W. Kolhörster.

Mit 8 Abbildungen. (Eingegangen am 18. Juni 1929.)

the nature of the „high-altitude radiation“

coincidence technique




Fig. 2.

absorber




Fig. 1.

GM tube

coinc./min




Fig. 3.

absorber thickness



W. Bothe
Nobel Prize 1954

W. Bothe & W. Kolhörster, Z. f. Phys. 56 (1929) 751

ist die Höhenstrahlung

the nature of the „high-altitude radiation“

Dreifachkoinzidenzen der Ultrastrahlung aus vertikaler Richtung in der Stratosphäre *).

I. Meßmethode und Ergebnisse.

Von Georg Pfozter in Stuttgart.

Mit 11 Abbildungen. (Eingegangen am 9. Juni 1936.)

Mit einer selbstaufzeichnenden Apparatur werden bei drei Registrierballon aufstiegen Dreifachkoinzidenzen der Ultrastrahlung aus vertikaler Richtung bis zu 10 mm Hg Luftdruck (29 km Höhe ü. M.) gemessen. Die Kurve der Zählrohrkoinzidenzen in Abhängigkeit vom Luftdruck zeigt ein Maximum bei 80 mm Hg und einen Buckel bei 300 mm Hg. Die Kurve kann gegen das Ende der Atmosphäre extrapoliert werden.





Fig. 6. Außen der Registrierapparatur. a) Von der Seite b) von oben gesehen.



Fig. 5. Launching of a balloon train from the courtyard of the institute.




Fig. 4. a) Aufstellplatte (nach Größe, Fülle); b) Vergrößerter Ausschnitt.

3) Die Firma Gebr. Junghans, Schramberg, hat uns freundlicherweise diese schönen Zählwerke hergestellt.

Dreifachkoinzidenzen der Ultrastrahlung aus vertikaler Richtung in der Stratosphäre *).

I. Meßmethode und Ergebnisse.

Von Georg Pfozter in Stuttgart.

of coincidences in 4 min




Fig. 1. Aus drei Aufstiegen gemittelte Kurve der Vertikalintensität der Ultrastrahlung in der Atmosphäre.

Latitude effect

Letters to the Editor

[The Editor does not hold himself responsible for opinions expressed by his correspondents. Neither can he undertake to return, nor to correspond with the writers of, rejected manuscripts intended for this or any other part of NATURE. No notice is taken of anonymous communications.]

Latitude Effect of Cosmic Radiation

ON the expedition organised by the Deutscher und Oesterreichischer Alpenverein in 1932 to the Andes of Peru, observations of cosmic rays were made at several heights up to 6,100m. and during the sea-voyage. From Bremen to Peru one apparatus worked during March and April 1932 on board the M.S. *Erfurt* of the Norddeutscher Lloyd line. On the return voyage in January and February 1933, three apparatuses were in full action from Peru through the Strait of Magellan to Hamburg on board the M.S. *Isis* of the Hamburg-Amerika line. The self-recording electrometers were constructed by Prof. E. Regener on the same principle as those used for his researches in Lake Constance¹ and in the upper atmosphere². The electrometer wire is inside an ionisation chamber of 16 cm. diameter with 'deltametal' walls of 1 cm. thickness. The position of the wire is photographed every half-hour on a fixed photographic plate.

Instrument No. 1 was filled with carbon dioxide at 9.7 atmospheres pressure and 16° C. With a radium capsule, I found the temperature effect on ionisation to be + 0.13 per cent for every + 1° C. difference. The correction for barometric pressure was 0.29 per cent per millimetre of mercury. All data were reduced to 16° C. and 760 mm. pressure. The ionisation due to radioactivity in the chamber itself was allowed for as 0.8 volts per hour as found on the bottom of Lake Constance at a depth of 250 m. Eight hemispherical shells of iron were fitted round the chamber. The combined thickness of this iron wall was 10 cm.

In Fig. 1 are recorded the data of apparatus No. 1, the iron case of which was open on the upper side. The graph shows the intensity of cosmic radiation in volts per hour for different geomagnetic latitudes on the voyage from the Strait of Magellan to Hamburg. The geographical position of the geomagnetic north pole was taken to be 78° 32' N. and 69° 08' W. Each point of the curve corresponds to an average of a twenty hours' registration. The points give a smooth curve which shows the accuracy of the recording method employed. The intensity increases by about 12 per cent when going from the equatorial region to 55° N. geomagnetic latitude.

Apparatus No. 2 was wholly encased in the iron shell. Apparatus No. 3 worked without any iron shell. Every instrument shows substantially the same effect.

In general, the curves agree with the observations of Clay³ and with those of A. H. Compton⁴ made at about the same time. It is very interesting that the northern and southern parts of the curve are not

symmetrical with respect to either the geomagnetic or the geographical equator. Considering the accuracy of our uninterrupted registration, this result is quite trustworthy.

From the fact that a latitude effect of 12 per cent of the radiation exists, it must be concluded that this part of the radiation consists of corpuscles before entering the earth's atmosphere. For the magnitude of this part of the radiation, reference should be made to the analysis of the components of cosmic rays by Regener² and Lenz⁵.

A more detailed report of these observations and of the researches in the Andes will be published in the *Zeitschrift für Physik*.

H. HOERLIN.

Physikalisches Institut
der Technischen Hochschule,
Stuttgart. June 8.

- ¹ Regener, E., *Z. Phys.*, **74**, 433; 1932.
- ² Regener, E., *Phys. Z.*, **34**, 306; 1933.
- ³ Clay, J., *Naturwiss.*, **20**, 687; 1932.
- ⁴ Compton, A. H., *Phys. Rev.*, **43**, 387; 1933.
- ⁵ Lenz, E., *Z. Phys.*; in the press.




FIG. 1.

Clay: Latitude Effect


RESULTS OF THE DUTCH COSMIC RAY EXPEDITION 1933

II. THE MAGNETIC LATITUDE EFFECT OF COSMIC RAYS
A MAGNETIC LONGITUDE EFFECT

by J. CLAY, P. M. VAN ALPHEN and C. G. 'T HOOFT

Natuurkundig Laboratorium, Amsterdam

journey from Holland to Java
intensity varies with latitude



cosmic rays are
charged particles




Fig. 1. Records of the variation of Cosmic Radiation with latitude on two different routes under different shielding with different instruments

- × × results with instrument *D* open (Amsterdam—Batavia)
- (*L*₁, *L*₂, *L*₃, *L*₄) results with instrument *D*₁ open (Batavia—Amsterdam)
- • Results 1928 and 1929.

Compton: World-wide survey of intensity of radiation




Fig. 24. The instrument used in this survey is usually shielded with lead and is placed in the box when used in most airplane flights.




Fig. 25. The camera will take a one-hundred-foot reel of 35 mm motion picture film which is driven at a constant rate past the slit by a power clock. Changeable gears allow various rates of film speeds to be used, depending on the expected ionization.




Fig. 27. Showing the type of record obtained at sea level in this world survey. Two of the horizontal lines give barometric and temperature records.

~1930

THE PHYSICAL REVIEW

A Journal of Experimental and Theoretical Physics

Vol. 43, No. 6

MARCH 15, 1933

SECOND SERIES

A Geographic Study of Cosmic Rays

ARVID H. COHEN, *University of Chicago*

(Received January 30, 1933)




FIG. 7. Intensity vs. geomagnetic latitude at sea level, including data of Clay and Millikan.




FIG. 6. Intensity vs. geomagnetic latitude for different elevations.

1931-34 A.H. Compton 12 expeditions → ~100 locations




FIG. 6.—Compton's world map of isocosms. Note the parallelism of these lines of equal cosmic-ray intensity and the dotted curves of geomagnetic latitude (50° N. and S.).



cosmic rays are charged particles

~1937 East-West Effect of Cosmic-Ray Intensity




Fig. 14. The equipment for the E-W experiment.

Rossi and others

higher intensity from the west

➡ cosmic rays are mostly positively charged

~1930 „elementary particles“: charged neutral

Rutherford (1919) **p** **n** (1932) Chadwick

Thomson (1897) **e⁻** **γ** (1905/26) Einstein



Discovery of new particles in cosmic rays

~1930 – 1950

birth of elementary particle physics



cloud chamber
C.T.R. Wilson
Nobel Prize 1927

