



Radboud Universiteit Nijmegen  
Jörg R. Hörandel and Sascha Caron  
Bjarni Pont

Particles and the Cosmos – 2019/20  
Werkcollege 11 – Cosmic ray propagation  
28.11.2019

**Problem 20** Spectra of cosmic-ray nuclei

The energy spectra of cosmic rays observed above the atmosphere are usually described by power laws

$$\frac{dN}{dE} \propto E^\gamma$$

at high energies (where solar modulation can be neglected). Compare the energy spectra for Carbon and Boron nuclei.

- Which one is steeper at high energies?
- Why?
- What is the conceptual difference between elements like carbon and elements like boron?

**Problem 21** Energy spectra of electrons

The energy spectrum of electrons in cosmic rays can be described by a power law with a spectral index of  $\gamma \approx -3.3$ .

- Why is the energy spectrum of electrons steeper as compared to nuclei (typical values  $\gamma \approx -2.7$ )?
- Enumerate the processes which occur during the propagation of electrons through the Galaxy.
- Calculate the synchrotron energy losses for an electron with an energy of 100 GeV in the Galaxy and compare it to the losses by protons. Consider a Galactic magnetic field strength of  $B = 3 \mu\text{G}$ . Calculate the power radiated as synchrotron radiation for both particle species and give the result in [eV/s]. Hint: the radiated power for a particle with charge  $e$  and energy  $E$ , moving on a circular trajectory with radius  $r$ , amounts to


$$P(E, r) = \frac{e^2 c}{6\pi\epsilon_0 r^2} \left( \frac{E}{m_o c^2} \right)^4 .$$

The permittivity of vacuum is given as  $\epsilon_0 = 8.85 \cdot 10^{-12} \text{ A s/V m}$ .

Estimate the time needed until the electrons and protons have radiated their complete energy through synchrotron radiation.

**Problem 22** Abundance of elements

The figure shows the abundance of elements in cosmic rays compared to the typical abundance in the solar system.



taken from J.R. Hörandel, *Advances in Space Research* 41 (2008) 442

Explain the main features of this diagram:

- p and He are less abundant - why?
- up and down of elements with even and odd charge numbers
- why are Li, Be, B more abundant in cosmic rays?
- why are the sub-iron elements more abundant in cosmic rays?

**problem 23** Muons in the atmosphere

In the lecture we discussed the production of pions in the atmosphere. The number of pions in the atmosphere reaches a maximum at an atmospheric depth around 100 to 200 g/cm<sup>2</sup>. Some of these pions decay into muons, i.e. the muons are produced at large heights in the atmosphere. The lifetime of a muon is  $\tau = 2.2 \mu\text{s}$ .

- How far can a muon travel before it decays?
- Why do muons travel down to sea level in the atmosphere?
- Calculate the decay length of a muon with an energy of 10 GeV

Student assistant: Bjarni Pont b.pont at astro.ru.nl

Lecture web site: <http://particle.astro.ru.nl/goto.html?partcos1920>