



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

Advances in Space Research 48 (2011) 1295–1303

**ADVANCES IN  
SPACE  
RESEARCH**

(a COSPAR publication)

[www.elsevier.com/locate/asr](http://www.elsevier.com/locate/asr)

## Thunderstorm observations by air-shower radio antenna arrays

W.D. Apel<sup>a</sup>, J.C. Arteaga<sup>b,1</sup>, L. Bähren<sup>c</sup>, K. Bekk<sup>a</sup>, M. Bertaina<sup>d</sup>, P.L. Biermann<sup>e</sup>,  
 J. Blümer<sup>a,b</sup>, H. Bozdog<sup>a</sup>, I.M. Brancus<sup>f</sup>, P. Buchholz<sup>g</sup>, S. Buitink<sup>c,2</sup>, E. Cantoni<sup>d,h</sup>,  
 A. Chiavassa<sup>d</sup>, K. Daumiller<sup>a</sup>, V. de Souza<sup>b,3</sup>, F. Di Pierro<sup>d</sup>, P. Doll<sup>a</sup>, M. Ender<sup>b,\*</sup>,  
 R. Engel<sup>a</sup>, H. Falcke<sup>c,i</sup>, M. Finger<sup>a</sup>, D. Fuhrmann<sup>j</sup>, H. Gemmeke<sup>k</sup>, C. Grupen<sup>g</sup>,  
 A. Haungs<sup>a</sup>, D. Heck<sup>a</sup>, J.R. Hörandel<sup>c</sup>, A. Horneffer<sup>e</sup>, D. Huber<sup>b,\*</sup>, T. Huege<sup>a</sup>, P.G. Isar<sup>a,4</sup>,  
 K.-H. Kampert<sup>j</sup>, D. Kang<sup>b</sup>, O. Krömer<sup>k</sup>, J. Kuijpers<sup>c</sup>, K. Link<sup>b</sup>, P. Łuczak<sup>l</sup>, M. Ludwig<sup>b</sup>,  
 H.J. Mathes<sup>a</sup>, M. Melissas<sup>b</sup>, C. Morello<sup>h</sup>, S. Nehls<sup>a</sup>, J. Oehlschläger<sup>a</sup>, N. Palmieri<sup>b</sup>,  
 T. Pierog<sup>a</sup>, J. Rautenberg<sup>j</sup>, H. Rebel<sup>a</sup>, M. Roth<sup>a</sup>, C. Rühle<sup>k</sup>, A. Saftoiu<sup>f</sup>, H. Schieler<sup>a</sup>,  
 A. Schmidt<sup>k</sup>, F.G. Schröder<sup>a</sup>, O. Sima<sup>m</sup>, G. Toma<sup>f</sup>, G.C. Trinchero<sup>h</sup>, A. Weindl<sup>a</sup>,  
 J. Wochele<sup>a</sup>, M. Wommer<sup>a</sup>, J. Zabierowski<sup>l</sup>, J.A. Zensus<sup>e</sup>

<sup>a</sup> Karlsruhe Institute of Technology (KIT), Institut für Kernphysik, Karlsruhe, Germany<sup>b</sup> Karlsruhe Institute of Technology (KIT), Institut für Experimentelle Kernphysik, Karlsruhe, Germany<sup>c</sup> Radboud University Nijmegen, Department of Astrophysics, The Netherlands<sup>d</sup> Dipartimento di Fisica Generale dell' Università Torino, Italy<sup>e</sup> Max-Planck-Institut für Radioastronomie Bonn, Germany<sup>f</sup> National Institute of Physics and Nuclear Engineering, Bucharest, Romania<sup>g</sup> Universität Siegen, Fachbereich Physik, Germany<sup>h</sup> INAF Torino, Istituto di Fisica dello Spazio Interplanetario, Italy<sup>i</sup> ASTRON, Dwingeloo, The Netherlands<sup>j</sup> Universität Wuppertal, Fachbereich Physik, Germany<sup>k</sup> Karlsruhe Institute of Technology (KIT), Institut für Prozessdatenverarbeitung und Elektronik, Karlsruhe, Germany<sup>l</sup> Soltan Institute for Nuclear Studies, Lodz, Poland<sup>m</sup> University of Bucharest, Department of Physics, Bucharest, Romania

Received 18 February 2011; received in revised form 31 May 2011; accepted 5 June 2011

Available online 12 June 2011

### Abstract

Relativistic, charged particles present in extensive air showers (EAS) lead to a coherent emission of radio pulses which are measured to identify the shower initiating high-energy cosmic rays. Especially during thunderstorms, there are additional strong electric fields in the atmosphere, which can lead to further multiplication and acceleration of the charged particles and thus have influence on the form and strength of the radio emission. For a reliable energy reconstruction of the primary cosmic ray by means of the measured radio signal it is very important to understand how electric fields affect the radio emission. In addition, lightning strikes are a prominent source of

\* Corresponding authors. Present address: KIT, Institut für Werkstoffe der Elektrotechnik, Karlsruhe, Germany (M. Ender).

E-mail addresses: [Moses.Ender@kit.edu](mailto:Moses.Ender@kit.edu) (M. Ender), [Daniel.Huber@kit.edu](mailto:Daniel.Huber@kit.edu) (D. Huber).

<sup>1</sup> Now at: Universidad Michoacana, Morelia, Mexico.

<sup>2</sup> Now at: Lawrence Berkeley National Laboratory, United States.

<sup>3</sup> Now at: Universidade de São Paulo, Instituto de Física de São Carlos, Brazil.

<sup>4</sup> Now at: ISS, Bucharest, Romania.

broadband radio emissions that are visible over very long distances. This, on the one hand, causes difficulties in the detection of the much lower signal of the air shower. On the other hand the recorded signals can be used to study features of the lightning development. The detection of cosmic rays via the radio emission and the influence of strong electric fields on this detection technique is investigated with the LOPES experiment in Karlsruhe, Germany. The important question if a lightning is initiated by the high electron density given at the maximum of a high-energy cosmic-ray air shower is also investigated, but could not be answered by LOPES. But, these investigations exhibit the capabilities of EAS radio antenna arrays for lightning studies. We report about the studies of LOPES measured radio signals of air showers taken during thunderstorms and give a short outlook to new measurements dedicated to search for correlations of lightning and cosmic rays.

© 2011 COSPAR. Published by Elsevier Ltd. All rights reserved.

*Keywords:* Cosmic rays; Radio detection; Thunderstorms; Lightning

## 1. Introduction

For studying high-energy cosmic rays above  $\approx 10^{15}$  eV ground based observatories covering large areas are needed because the flux of the cosmic rays gets too low at these energies for direct measurements by balloon or satellite bound experiments. By ground based detection arrays instead, the secondary particles of an extensive air shower (EAS) generated by the impinging primary cosmic ray are measured. Beside the standard method of EAS detection with an array of particle detectors, one possibility to observe these secondary particles is to measure the radio emission that is produced by the electrons and positrons due to their deflection in the Earth's magnetic field. This radio emission gives an integrated measurement all over the shower development in the atmosphere contrary to particle detectors that only record a footprint of the shower at observation level. In contrast to the fluorescence or Cherenkov detection technique, where also the longitudinal shower development is observed by means of the EAS's electromagnetic emission, radio signals can be measured at night and daytime (Haungs et al., 2003). The radio detection technique can complement the information gained by particle detectors as well as can be operated as a stand alone detection technique. The problem of this technique lies in the weak signal compared to the ambient noise. But, with highly sensitive radio antenna arrays most of the noise sources can be identified and the data corrected for Schröder et al. (submitted for publication), except one of the most disturbing noise source formed by the radio emission of lightning strikes.

Lightnings have been explored for a very long time, but the mechanism that leads to the discharging process is still not fully understood (Dwyer et al., 2008). There are several theories that could explain the sudden electric breakdown, where a very intriguing theory predicts that lightning strikes might be initiated by cosmic rays. To validate this theory, cosmic rays and lightning strikes need to be measured at the same place and time. As lightning observations are most effective in radio or X-ray due to the given transparency of the thunderclouds in these wavelengths, it seems natural to use EAS radio antenna arrays for such measurements.

The digital radio antenna array LOPES (Falcke et al., 2005; Haungs et al., 2009) was designed and built to

perform the proof-of-principle of the EAS radio detection technique. LOPES is co-located and combined with the air-shower detector KASCADE-Grande (Antoni et al., 2003; Apel et al., 2010). KASCADE-Grande is a multi-detector air-shower experiment located in Karlsruhe, Germany, and consists of different particle-detector devices to measure all kinds of secondary particles in a large primary energy range of  $10^{14}$ – $10^{18}$  eV.

In order to study the effect of large atmospheric electric fields on the EAS radio signal present during thunderstorms, detailed investigations are performed at LOPES. It is shown that LOPES is able to detect air showers and, in addition, lightning strikes. Hence, LOPES can be used as a lightning mapping array and the correlation between EAS and lightning strikes is studied by getting the position and time of the lightning strike from LOPES and the air shower data from KASCADE-Grande. So far no correlation is seen, but with the technology used at LOPES the quality of the information as a lightning mapping array is rather low. However, by deploying lightning mapping systems that are particularly designed for lightning detection, see (Rison et al., 1999), e.g., a far better time and spatial resolution of the lightning can be achieved. In addition, an increase of the lightning statistics would be desirable as the area covered by LOPES and KASCADE-Grande is small compared to the size of thunderstorm clouds. A future project will combine all these by deploying special designed lightning detectors at a large area cosmic ray surface experiment. By covering a huge area and having high quality information of both, lightning strikes and cosmic rays, one will be able to decide whether there is a correlation between lightning strikes and cosmic rays or not.

In this article we summarize the activities within LOPES with respect to thunderstorm observations and briefly discuss the prospects of future projects.

## 2. EAS observations by radio antenna arrays

The measurement of cosmic ray induced air showers can be performed with different detection techniques. One of these techniques is the analysis of the radio pulses emitted by the electromagnetic component of an air shower. Using this method a complementary measurement to the settled technique of particle detectors is achieved. Several radio

antenna arrays have been built or are going to be constructed, where LOPES in Karlsruhe served as the ground breaker for this cosmic ray detection technique. The Auger Engineering Radio Array AERA at the Pierre Auger Observatory in Argentina, presently being constructed, is the first array of a large scale application of the technique (van den Berg et al., 2009).

By an interferometric standard analysis of a radio pulse measured by a dedicated radio antenna array several shower parameters can be determined, e.g. the arrival direction and the energy of the primary particle, or the shower maximum and therefore the mass of the incoming cosmic ray. To reconstruct these quantities several steps have to be performed in such an analysis, where in the case of LOPES this is done as follows: to link the measured raw ADC counts with the field strength of the radio pulse the attenuation of the complete signal chain is investigated using an external source to calibrate the signal. The raw ADC counts for individual events are corrected for the electronic delay, attenuation, and dispersion by shifting the time traces and by multiplying the calibration factors. After that the narrow band noise is suppressed, where for this procedure the data are Fourier transformed from the time to the frequency domain. Although the pulses are filtered to a bandwidth where no broadcasting stations are operating in Karlsruhe, i.e. 40–80 MHz, several narrow band noise sources can be seen in the spectra. Then, the reconstruction begins using the arrival direction of the cosmic ray measured by KASCADE-Grande as starting value. Taking into account the direction sensitive antenna characteristics, a so called beam forming is done. Such a beam forming is an iterative procedure to find the coherent part of the received signal in many antennas by shifting artificially the times until the direction is found where the coherent signal has its maximum. The beam forming in case of LOPES is based on the calculation of the cross-correlation value, which has a higher sensitivity to the coherent part of the signal than, e.g., a simple power beam. Calculating the cross-correlation beam leads to an increase of the signal-to-noise ratio, because the coherent part of a signal is amplified and the incoherent part, the noise, is suppressed. This way the entire antenna array is used and even EAS signals that are not seen in individual antennas are reconstructed. Since LOPES is sampling in the second Nyquist domain no information is lost, though the signal is recorded with a lower sampling rate than the highest frequency of the bandwidth to keep the recorded data small and save disk space. By an up-sampling procedure the original information is recovered.

Fig. 1 shows the time traces of all 30 LOPES antennas and the corresponding cross-correlation beam for an individual radio detected cosmic ray event. The time distribution of the cross-correlation beam values (top panel of Fig. 1) is fitted with a Gaussian function in order to obtain the radio EAS observables, where the most important is the field strength value at the maximum of the Gaussian function (bottom panel of Fig. 1). This field strength value is directly correlated with the energy of the primary cosmic




Fig. 1. Example of an EAS detection by the LOPES antenna array. Top: bandwidth normalized field strength in  $\mu\text{V/m/MHz}$  vs. time in  $\mu\text{s}$ . The traces of all 30 LOPES channels are shown. Bottom: cross-correlation beam and Gaussian fit. The cross-correlation beam assigns time dependence of the averaged coherent part of the radio power received from the extensive air shower.

ray, only corrected for the geometry of the shower, i.e. arrival direction and position of incidence on observation level with respect to the antenna positions.

So far, LOPES has detected more than a thousand high-energy cosmic ray events serving as input on detailed analysis of the radio signal and the understanding of the radio emission in air showers in order to pave the way for large scale applications of the technique (Haungs et al., 2009).

### 3. Atmospheric electric field during thunderstorms

The exact measurement of the atmospheric electric field with a high time resolution is very important to provide information for the radio detection of cosmic ray induced air showers. The radiation in the radio regime is emitted by the propagating electrons and positrons of an air shower mainly due to their time dependent spatial (charge) separation in the Earth's magnetic field. The atmospheric electric field has an influence on the propagation of the charged particles of the shower and therefore on the radio emission. Only knowledge of the field allows detailed studies of the influence of the atmospheric electric field on the radio emission.

To measure the atmospheric electric field a field mill is used. With such a device it is possible to record the vertical electrical field between the lowest cloud layer and the ground. The electric field strength gives clear evidences whether there are fair weather conditions or a close thunderstorm.

During fair weather conditions, the atmospheric electric field experiences only small changes between  $-100$  and  $-200 \text{ Vm}^{-1}$ . The amplitudes increase when rain clouds cross over but the changes are on large time scales. During thunderstorms the field strength can reach values up to  $\pm 20 \text{ kVm}^{-1}$  and there are sudden changes in the electric field occurring on a very short time scale, see Fig. 2.

#### 4. Influence of atmospheric electric fields on radio detection of EAS

An additional or strongly varying atmospheric electric field can lead to different strengths and geometries of the radio emission from charged particles, in particular secondaries from a cosmic-ray air shower in the Earth's atmosphere (Buitink, 2007; Buitink et al., 2010). In Fig. 3, this effect is schematically shown. The electric field in thunderclouds, especially within the convective region, can reach values up to  $\pm 100 \text{ kVm}^{-1}$ . With radio antenna arrays, like




Fig. 3. Scheme of the influence of an additional electric field on the electrons and positrons of an air shower. On the left hand side no additional electric field is shown. In the middle, an electric field parallel to the direction of the air shower is shown which leads to an acceleration of the positrons and a deceleration of the electrons. On the right hand side, an electric field which is perpendicular to the shower direction is shown which results in a stronger deflection of the electrons and positrons (from Ref. Buitink, 2007).

e.g. LOPES, this effect can be seen in the observation of cosmic rays by recording an amplified or weakened radio signal during thunderstorms and extreme weather conditions. In order to detect a thunderstorm a field mill has




Fig. 2. Electric field measured by an E-field mill. Top: during fair weather conditions. Bottom: during a thunderstorm, where discontinuities and jumps in the electric field strength are seen. In addition, a much higher field strength than during fair weather is measured.




Fig. 4. Bandwidth normalized field strength in  $\mu\text{V/m/MHz}$  vs. time in  $\mu\text{s}$  for two events with similar geometry and primary energy. Top: time traces of an event recorded during fair weather conditions, where no coherent signal can be seen. The increase in some of the traces starting at  $-1.75 \mu\text{s}$  is assigned to detector noise from the KASCADE particle detectors. Bottom: time traces of a similar event which was recorded during a thunderstorm. The amplified coherent signal from the direction of the air shower at  $-1.8 \mu\text{s}$  is clearly visible.

been installed at the LOPES site (Nehls, 2008) serving as a monitoring and veto device for the LOPES analysis. One example for such an amplification can be seen in Fig. 4 where two events with very similar shower geometry and primary energy are shown. The first event was recorded during fair weather conditions with no observable radio signal in the trace which should occur at about  $-1.8 \mu\text{s}$ . This is expected for the low estimated primary energy by KASCADE-Grande observations of  $5.4 \times 10^{16}$  eV. The incoherent signal starting at  $-1.75 \mu\text{s}$  is assigned to detector noise from the KASCADE particle detectors. This air shower arrived at the KASCADE array with  $\phi = 110.39^\circ$  and  $\theta = 31.5^\circ$ , where  $\phi$  is the KASCADE reconstructed azimuth of the shower direction and  $\theta$  the zenith angle. The second event is an air shower with  $\phi = 110.35^\circ$  and  $\theta = 32.1^\circ$  and an even lower energy estimated to  $4.3 \times 10^{16}$  eV. The average distance of the antennas to the shower core is also in the same order for both events. In the time trace of this event a coherent radio pulse at  $-1.8 \mu\text{s}$  is observed, but for that energy no radio signal is expected. Most likely explanation for such a clear detection is the amplification of the EAS radio signal in the strong electric fields of the thundercloud present at that time.

The energy threshold for triggering LOPES by KASCADE-Grande is lower than the possible detection threshold ( $5 \times 10^{16}$  eV) at the site of LOPES with its industrial environment and high noise level. Because of that only in a small fraction of the triggered events a radio pulse can be observed. In a fraction of  $(0.96 \pm 0.12) \times 10^{-2}$  of the events recorded during fair weather conditions a coherent radio signal has been seen. In 2007 and 2008 approximately two full days of data were recorded at periods of thunderstorms. Because of these low statistics only few such corresponding fair weather – thunderstorm partner events could be analyzed. But the fraction of events that were recorded during thunderstorms and that show a coherent signal is with  $(2.39 \pm 0.27) \times 10^{-2}$  about a factor two to three higher than in fair weather conditions. This results in 81 events that are recorded during thunderstorms having a cross-correlation beam above threshold. It serves as a clear indication that strong atmospheric electric fields during thunderstorms have an influence on the radio emission of cosmic ray air showers and might more likely amplify the signal than attenuate it. For smaller atmospheric electrical fields there might be no big effect (Buitink, 2007) but recent investigations have shown that strong atmospheric electric fields without thunderstorm that can occur in rain clouds also have an influence on the radio emission but the number of events recorded during such conditions are even lower (Ender et al., 2009).

The aim of such studies is that by knowing the influence of electric fields on the radio emission in EAS the radio signal recorded during extreme weather conditions can be corrected for and therefore these periods are not lost for the analysis. This could lead to an uptime for the radio detection technique close to 100%, which is important as the flux of high-energy cosmic rays is very low. For the time being

the atmospheric monitoring is used as veto for the standard operation of radio antenna arrays such as LOPES.

## 5. Radio background during thunderstorms

To achieve the desired timing accuracy for beam forming reconstruction a reference antenna (beacon) was installed at LOPES. The beacon emits constant sine waves at 63.5, 68.1 MHz and (since end of 2010) 53.1 MHz which form a considerable background of the measurements during fair weather. During thunderstorms the general ambient background is much higher and in addition radio signal from lightning strikes contribute to the background. In Fig. 5 two average background frequency spectra are shown. The upper spectrum was recorded during fair weather conditions. The narrow band noise and the peaks of the beacon signals are clearly seen. The lower part of the figure shows a spectrum taken during thunderstorm conditions. Here, the beacon signals and other narrow band noise sources can hardly be identified over the high broadband radio emission. As the beacon signals are overlapped in such periods the recorded events cannot be analyzed.




Fig. 5. Logarithm of the mean amplitude of background frequency spectra in arbitrary units. Top: background spectrum recorded during fair weather conditions, where a lot of narrow band noise sources are observed as small peaks. Bottom: background spectrum recorded during a thunderstorm. The much higher broad band background level exceeds nearly all narrow band noise peaks including the beacon signals needed for the event-to-event time calibration.

Another aspect that makes it impossible to analyze individual events are the fact that the sensitive electronics needed to observe the weak radio signals from cosmic-ray air showers are saturated by the strong signals from lightnings.

To be competitive with other EAS detection techniques it is aspired to trigger on the radio pulse only, in order to be independent from external triggers, e.g. by particle detectors. Within LOPES<sup>STAR</sup> (Asch, 2009) such a self-trigger technique is under development. The trigger condition is based beside other conditions on the pulse shape. To avoid wrong trigger decisions that arise from thunderstorms it is important to know the characteristics of non-EAS short pulses, e.g. generated by lightning.

## 6. Problem of lightning initiation

One of the most extreme weather conditions mankind can think of are thunderstorms and lightning. Although these appearances are known for a very long time and have been the focus of many studies, the mechanism that leads to the final electric breakdown is still not well known. The field strengths of the electric fields in thunderclouds are large, but too small for a classical breakdown. One mechanism that could explain how a breakdown can happen with smaller electric field strengths are relativistic runaway electron avalanches (RREA). These RREA occur at a certain critical field strength when the cross-section of the electron-electron interaction gets smaller which leads to ionization losses that are smaller compared to the energy gain at the critical field strength. This results in a net energy gain of the electrons and an increasing number of electrons. This is since the electrons coming from the ionization are also accelerated and again produce new unbound electrons. This mechanism can only take place when the number of unbound electrons that can be accelerated, the seed electrons, is high enough within the strong electric field of a thundercloud. A source that can provide these seed electrons are cosmic ray induced air showers. During an air shower development up to  $10^6$  electrons and positrons can be generated on a very limited area, which is the location of the shower maximum, typically in a height of 3–8 km above sea level. These particles are then accelerated in the strong electric field of the thundercloud and lead to a RREA which results in a breakdown, called RRB, relativistic runaway breakdown. So, cosmic rays could be the initiator for lightning by providing the seed electrons for a RREA (Dwyer, 2009).

## 7. Radio signal from lightning strikes

The jumps and discontinuities in the electric field are clear evidences for thunderstorms and are used at LOPES to change the data acquisition into the so-named thunderstorm mode (Nehls, 2008). During this special mode the recorded time traces are roughly eight times longer than the usual 0.82 ms and 6.55 ms of data taken for each trig-

gered event, where the pre-trigger time of 0.41 ms remains the same. This is done to be able to look for timely extended lightning signals in the recorded traces visible after the EAS signal, see an example in Fig. 6. A discharging process like a lightning is always accompanied by strong electric fields in thunderclouds and always emits broadband electromagnetic radiation (Rakov and Uma, 2005). The radiation can be observed in the radio regime over long distances with antenna arrays originally designed to detect the radio emission from cosmic-ray air showers. The time structure of the signal can be very different depending on the distance and the nature of the discharging process. Fig. 6 shows a lightning signal recorded by all antennas of the LOPES array.

Discharges located in clouds with altitudes between 5 and 20 km above sea level can produce very short and strong pulses that are described as narrow bipolar pulses by Gurevich and Zybin (2004). It could be an interesting question to investigate if such short pulses seen in Fig. 6 just in front of the large emission from the lightning strike are generated by relativistic runaway breakdowns (RRB) and if such RRB's are always present at the initiation of lightning strikes.

Discharges within or between clouds or from cloud to ground result in signals that are lasting longer than RRBs or EAS signals (see Fig. 6). This gives us the possibility to calculate for many successive time slots and in every direction of the sky the cc-beam value and to combine them into a skymap. Typically the cross-correlation beam is used, since the signal from lightning strikes is coherent, but if desired also the power beam (averaged power of all antennas in a certain direction) is used to generate the skymap. By this procedure the lightning development is observed and the location of origin, the type and the direction of the lightning is reconstructed. For the principles of lightning observation using radio interferometers see also




Fig. 6. Example for time traces of the 30 LOPES antennas recorded in the thunderstorm mode for a KASCADE triggered air shower. Whereas the cosmic ray event triggering the readout is visible as small peak in the beginning of the traces, in this particular event a lightning strike occurs approximately 4 ms after the EAS. Structures in the radio emission of this lightning are very nicely resolved in all 30 antennas.




Fig. 7. This figure shows a skymap of a lightning recorded with LOPES, where for each cell the amplitude of the cross-correlation beam is displayed. The whole sky is shown with the zenith in the center of the plot and the horizon at the edge. The lightning can be seen as the strong signal in the North–Northwest (circle). The signals spread over the whole sky are due to the grating lobes of the antenna array which is a well understood artifact of skymapping in astronomy and only dependent on the geometry of the antenna positions on ground. The event corresponds to the traces shown in Fig. 6 and the star assigns the location of the detected EAS in this event.

(Rhodes et al., 1994). For example the lightning strike displayed in Fig. 7 (which corresponds to the time traces shown in Fig. 6) is a cloud-to-cloud lightning as the track does not reach the horizon. This example shows the capabilities of EAS radio antenna arrays to investigate in detail the radio emission from individual lightning discharges.

## 8. Correlations of EAS and lightnings

To study the possibility of cosmic rays causing lightning strikes there are two correlations to investigate, the time and the spatial connections of air showers and lightning.

To investigate the timely correlation, at LOPES a lightning is detected by the E-field mill, where a jump in the electric field corresponds to a discharge process. If cosmic-ray air showers induce these lightning strikes they should be observed by KASCADE-Grande or LOPES shortly before the lightning. In Fig. 8, the time differences between jumps in the electric field with detected cosmic ray events is shown. The time binning of one second is determined by the time resolution of the electric field mill used at the LOPES site. In order to investigate systematically introduced uncertainties, similar plots with same dataset but artificially introduced time delays to scramble the possible correlation were produced. These studies resulted in only statistically insignificant enhancements at  $t_{diff} = 0$ .

The time resolution of the electric field mill is too low for such an analysis, but is high enough to determine whether




Fig. 8. Time correlation between lightning and air-shower events. Shown is the time difference between cosmic-ray event and lightning, where the lightning is detected by an electric field mill.

there is a thunderstorm or not. To see a time correlation between cosmic-ray air showers and lightnings, a better time resolution is necessary. This can be provided by LOPES, but the statistics for such an analysis is still too low at LOPES. In addition, the area where EAS can be detected is much smaller than the sensitive area for lightning strikes, which also worsen the search for time correlations.

A more promising correlation study between cosmic rays and lightning is the search for a combined spatial and time correlation. To study this it is essential to detect lightnings with a high spatial resolution and to detect cosmic-ray air showers with a high efficiency and also a good spatial resolution. For the lightning detection a LOPES like array can be used as a combined lightning mapping and EAS detection array. The direction of the lightning can be determined by calculating the cross-correlation beam in every direction of the sky. A calculated skymap is shown in Fig. 7, where an intracloud discharge can be seen. A star marks the direction of the air shower arriving shortly before the lightning. In this event and also in others no correlation could be observed. To improve the searching for spatial correlations a better detection of the lightning and especially the lightning development is needed. Not only the direction where the lightning happened is of interest but also the path of the lightning and whether a cosmic-ray air shower passed there at the start of the lightning or somewhere near. The path of the air shower can be observed very well and reconstructed with KASCADE-Grande and LOPES. The development of the lightning is difficult to reconstruct with the given instruments since LOPES was not designed for that kind of studies and still the covered area is too small.

## 9. Future investigations

The LOPES experiment, which will stop operation by 2011, experienced some changes in hardware, but still will be able to measure in the thunderstorm mode in the summer period 2011. Now, LOPES consists of 10 antenna

stations with each having three simple dipole antennas sensitive to the east–west, north–south, and vertical polarization direction, respectively. This makes the antenna array more sensitive, in particular to signals arriving from more inclined sources, i.e. from the horizon. In addition, electric field mills, an additional small lightning mapping array, and three antennas sensitive to radiation in the kHz frequency range will be active during the thunderstorm periods. In particular the kHz-information can be very useful for a combined multi-parameter observation of air-shower and lightning, as lightning strikes are strong emitters in these frequency domains. The KASCADE-Grande detector stations will be used to study the particle densities during a thunderstorm.

Among different possible techniques, radio observations with dedicated antenna arrays are most suitable to follow the lightning development in time and space with highest resolution. Such lightning mapping arrays (LMA) (Rison et al., 1999) are devices especially designed to better measure the initiation and generation of lightning and can be used – when co-located to air-shower experiments – to investigate the influence of thunderstorms and lightning on the air shower detection by particle detectors. The “Lightning Air Shower Study” project LASS as a part of the Pierre Auger Observatory (Abraham et al., 2004) is in this sense the next generation experiment for lightning and thunderstorm investigations in combination with an air-shower experiment. LASS will help to better understand the development of lightning as well as to investigate the influence of thunderstorms and lightning to the air shower detection by particle detectors, by fluorescence telescopes, and by radio antenna arrays such as LOPES or AERA, where AERA will be combined with the lightning mapping array.

By deploying different types of radio antennas and highly sensitive electric field mills a lightning can be observed with a spatial resolution of around 10 m and time resolution of about 40 ns (Thomas et al., 2004). Data from the LMA stations can then be combined to provide three-dimensional images of the lightning channels, including lightning initiation positions and times of occurrence. Each LMA station records peak signal magnitude and time in every 80  $\mu$ s interval in a quiet 6 MHz VHF band at typically 63 MHz.

The goals of such a project are the determination of a possible correlation between lightning and cosmic-ray air showers, to study the effects of strong electric fields on the radio emission of EAS, to study the influence of strong electric fields or thunderstorms on the particle component of EAS and to investigate whether distant lightnings harm the performance of the fluorescence detectors. For that purposes the installation within an existing EAS experiment, such as the Pierre Auger Observatory, is ideal. Both projects could profit from each other as lightning can also affect the air-shower measurements by producing a broadband light flash that can harm fluorescence measurements over long distances. The X- and gamma-rays

produced by lightning strikes can irradiate the particle detectors and cause background. The moving charge in a lightning causes also emission in the MHz range which causes additional radio background. The strong atmospheric electric fields during thunderstorms can seriously have an influence on the particle distributions of charged particles from an EAS in the atmosphere, which strongly affects the correct reconstruction of this air shower.

## 10. Conclusion

The LOPES experiment has provided the proof of principle for the radio detection technique of high-energy cosmic rays and has made key contributions to the understanding of the radio emission physics such as the predominantly geomagnetic origin of the emission. LOPES significantly contributes to the calibration of the shower radio emission in the primary energy range of  $10^{16}$ – $10^{18}$  eV. I.e., is investigating in detail the correlation of the measured field strength with the primary cosmic ray parameters, in particular the arrival direction, the energy and the mass. The radio detection technique has a high reliability in all but the most extreme weather conditions, as for example the reconstructed energy of an air shower is influenced by electric fields during thunderstorms, but not at normal weather conditions leading to a larger fraction of events with a detected coherent signal. Therefore it is mandatory for antenna air-shower arrays to monitor the electric field of the atmosphere.

Lightning strikes are one of the most extreme weather appearances that can occur in nature. Although they have been analyzed for a very long time, the basic mechanism that leads to the final breakdown is still not known. By observing lightning strikes in the radio regime the strikes can be observed looking through the thunderclouds since these are transparent for radio waves. This gives huge advantage to the studies of lightning in the radio regime over optical observations. Clouds are opaque for optical transmission which makes it impossible to observe lightning strikes properly. With the radio antenna array LOPES lightning strikes could be observed, although LOPES was not especially designed for that. LOPES was built within the existing cosmic-ray air shower experiment KASCADE-Grande. This allowed first low-level studies of correlations between cosmic rays and lightning strikes, by analyzing the air-shower properties given by KASCADE-Grande and the geometry of the lightning strike observed with LOPES. These first analyses showed no significant correlation between cosmic rays and lightning strikes.

The LOPES results are interesting and verified the capabilities of such arrays in lightning studies, but because of the drawbacks of a small area covered and short time traces recorded a final conclusion is not possible yet. To draw a final conclusion whether cosmic rays induce lightning strikes or not, more sophisticated studies with devices that are specially designed for that purpose need to be performed. This will be realized by future



experimental devices, such as LASS, where high tech air shower analyses and lightning detection over a huge area can be combined.

### Acknowledgment

LOPES and KASCADE-Grande have been supported by the German Federal Ministry of Education and Research. KASCADE-Grande is partly supported by the MIUR and INAF of Italy, the Polish Ministry of Science and Higher Education and by the Romanian Authority for Scientific Research CNCSIS-UEFISCSU (grant IDEI 1442/2008). Part of this research has been supported by grant number VH-NG-413 of the Helmholtz Association. D.H. and A.H. would like to thank Paul Krehbiel and his colleagues from New Mexico Tech, Joe Dwyer from Florida State University, and in particular Bill Brown, Colorado State University, for the initiation of the LASS project and the possibility to participate on these coming efforts. D.H. would like to thank the organizers of the TEPA conference held in Armenia, in particular Ashot Chilingarian, for the possibility to discuss the topic in an excellent atmosphere.

### References

- Abraham, J., Aglietta, M., Aguirre, I.C. et al. Properties and performance of the prototype instrument for the Pierre Auger observatory. *NIM A* 523, 50–95, doi:10.1016/j.nima.2003.12.012, 2004.
- Antoni, T., Apel, W.-D., Badea, F. et al. KASCADE Collaboration The cosmic-ray experiment KASCADE. *NIM A* 513, 490–510, 2003.
- Apel, W.D., Arteaga, J.C., Badea, F. et al. KASCADE-Grande Collaboration The KASCADE-Grande experiment. *NIM A* 620, 202–216, 2010.
- Asch, T. FZKA report 7459, Measuring the radio emission of cosmic ray air showers with LOPES, Forschungszentrum Karlsruhe, 2009.
- Buitink, S., Huege, T., Falcke, H., Kuijpers, J. Simulation of radio emission from air showers in atmospheric electric fields. *Astropart. Phys.* 33 (5–6), 296–306, 2010.
- Buitink, S., Apel, W.-D., Asch, T. et al. LOPES Collaboration Amplified radio emission from cosmic ray air showers in thunderstorms. *A&A* 467, 385–394, 2007.
- Dwyer, J.R., Uman, M.A., Rassoul, H.K. The remote measurement of thunderstorm electrostatic fields, *J. Geophys. Res.*, 114, D09208, 2008, doi:10.1029/2008JD011386.
- Dwyer J.R., Diffusion of relativistic runaway electrons and implications for lightning initiation, *J. Geophys. Res.*, 115, A00E14, 2009, doi:10.1029/2009JA014504.
- Ender, M., Apel, W.-D., Arteaga, J.C. et al. LOPES Collaboration, EAS radio emission during thunderstorms. In: Proceedings of the 31st ICRC, Lodz, Poland, 2009.
- Falcke, H., Apel, W.-D., Badea, F. et al. LOPES Collaboration Detection and imaging of atmospheric radio flashes from cosmic ray air showers. *Nature* 435, 313, 2005.
- Gurevich, A.V., Zybin, K.P. High energy cosmic ray particles and the most powerful discharges in thunderstorm atmosphere. *Phys. Lett. A* 329, 341–347, 2004.
- Haungs, A., Rebel, H., Roth, M. Energy spectrum and mass composition of high-energy cosmic rays. *Rep. Progress Phys.* 66, 1145, 2003.
- Haungs, A., Apel, W.-D., Arteaga, J.C. et al. LOPES Collaboration Air shower measurements with the LOPES Radio antenna array. *NIM A* 604, 1–8, 2009.
- Nehls, S. Calibrated measurements of the radio emission of cosmic ray air showers, FZKA report 7440, Forschungszentrum Karlsruhe, 2008.
- Rakov, V., Uma, M. *Lightning: Physics and Effects*. Cambridge Univ. Press, ISBN 0-521-03541-4, 2005.
- Rhodes, C.T., Shao, X.M., Krehbiel, P.R., Thomas, R.J., Hayenga, C.O. Observations of lightning phenomena using radio interferometry. *J. Geophys. Res.* 99, 13059–13082, 1994.
- Rison, W., Thomas, R.J., Krehbiel, P.R., Hamlin, T., Harlin, J. A GPS-based three-dimensional lightning mapping system: initial observations in central New Mexico. *Geophys. Res. Lett.* 26 (23), 3573–3576, 1999.
- Schröder F.G., Apel, W.-D., Arteaga, J.C. et al. LOPES Collaboration, On noise treatment in radio measurements of cosmic ray air showers. *Nucl. Instr. Meth. A – ARENA Proceedings*, submitted for publication, arXiv:1009.3444.
- Thomas, R.J., Krehbiel, P.R., Rison, W., Hunyady, S.J., Winn, W.P., Hamli, T., Harlin, J. Accuracy of the lightning mapping array. *J. Geophys. Res.* 109, D14207, doi:10.1029/2004/JD004549, 2004.
- van den Berg A., Abraham, J., Aglietta, M. et al. Pierre Auger Collaboration, Radio detection of high-energy cosmic rays at the Pierre Auger Observatory In: Proceedings of the 31st ICRC, Lodz, Poland, 2009.