

The origin of galactic cosmic rays

Jörg R. Hörandel

*Department of Astrophysics, Radboud University Nijmegen,
P.O. Box 9010, 6500 GL Nijmegen, The Netherlands
<http://particle.astro.kun.nl>*

The origin of galactic cosmic rays is one of the most interesting unsolved problems in astroparticle physics. Experimentally, the problem is attacked by a multi-disciplinary effort, namely by direct measurements of cosmic rays above the atmosphere, by air shower observations, and by the detection of TeV γ rays. Recent experimental results are presented and their implications on the contemporary understanding of the origin of galactic cosmic rays are discussed.

1 Introduction

The Earth is permanently exposed to a vast flux of highly energetic particles, fully ionized atomic nuclei from outer space. The extraterrestrial origin of these particles has been demonstrated by V. Hess in 1912¹ and he named the particles "Höhenstrahlung" or "Ultrastrahlung". In 1925 R. Millikan coined the term "Cosmic Rays". They have a threefold origin. Particles with energies below 100 MeV originate from the Sun. Cosmic rays in narrower sense are particles with energies from the 100 MeV domain up to energies beyond 10^{20} eV. Up to several 10 GeV the flux of the particles observed is modulated by the 11-year cycle of the heliospheric magnetic fields. Particles with energies below 10^{17} to 10^{18} eV are usually considered to be of galactic origin. A proton with an energy of 10^{18} eV has a Larmor radius $r_L = 360$ pc in the galactic magnetic field ($B \approx 3 \mu\text{G}$). This radius is comparable to the thickness of the galactic disc and illustrates that particles at the highest energies can not be magnetically bound to the Galaxy. Hence, they are considered of extragalactic origin. (A more detailed discussion of cosmic rays at the highest energies is given elsewhere^{2,3}.)

The energy density can be inferred from the measured differential energy spectrum⁴ dN/dE

$$\rho_E = \frac{4\pi}{c} \int \frac{E}{\beta} \frac{dN}{dE} dE, \quad (1)$$

where βc is the velocity of particles with energy E . For galactic cosmic rays the major contribution to the total energy density originates from particles with energies around 1 GeV. Such particles are strongly influenced by the heliospheric magnetic fields. Outside the heliosphere, in the local interstellar environment an energy density $\rho_E^{LIS} = 1.1 \text{ eV/cm}^3$ is obtained (Ref.⁵ p.12). The parameterization of the measured galactic cosmic-ray flux according to the polygono model⁶ results in a density $\rho_E^{gal} = 0.43 \text{ eV/cm}^3$. This is the measurable energy density at Earth (for an average modulation parameter $M = 750 \text{ MeV}$, see (1) in Ref.⁶). This implies that less than half of the energy flux can be registered directly at Earth.

In this article we will give an overview on recent experimental results and their implications on the contemporary understanding of the origin of galactic cosmic rays. As space is limited here, the reader may also consider further recent reviews by the author^{6,7,8,9,10,11}.

Progress in the understanding of the origin of galactic cosmic rays emerged mainly from observations in three complementary disciplines. The direct measurement of cosmic-ray particles above the atmosphere in outer space and on stratospheric balloons (Sect. 2). At energies exceeding 10^{15} eV the steeply falling cosmic-ray energy spectrum requires experiments with large detection areas exposed for long times, at present, only realized in ground based installations (Sect. 3). And the observation of TeV γ -rays (Sect. 4).

2 Direct measurements

Detectors above the atmosphere allow a precise determination of the composition of cosmic rays and the measurement of energy spectra for individual elements, presently, up to about 10^{14} eV. These findings are very important to understand the propagation of the particles in the Galaxy.

In the 100 MeV energy domain detailed studies of the isotopic composition of cosmic rays and the comparison to the local galactic abundance indicates that cosmic rays are accelerated out of a sample of well mixed interstellar matter¹². The propagation time of the particles in the Galaxy has been determined to be $\tau_{esc} = (15.6 \pm 1.6) \cdot 10^6$ yr^{13,14,15}. At GeV energies the elemental composition has been studied in detail, finding all elements of the periodic table in cosmic rays, see e.g.¹⁰. Some elements, as e.g. boron are produced during the propagation in the Galaxy through spallation of heavier elements such as carbon. Investigations of the abundance ratios of different elements (like the boron-to-carbon ratio) yield the column density of matter traversed by the particles in the Galaxy^{13,16}.

At present, several groups try to extend direct measurements to higher energies towards the knee (at about $4.5 \cdot 10^{15}$ eV) in the all-particle spectrum, benefiting from long-duration balloon flights¹⁷. Among these are the ATIC¹⁸, CREAM¹⁹, and TRACER²⁰ experiments. The latter is presently with an aperture of $5 \text{ m}^2 \text{ sr}$ the largest cosmic-ray detector for direct measurements. It comprises several electromagnetic detectors, namely an acrylic Čerenkov counter, a proportional tube array, and a transition radiation detector, thus covering a large energy range from 10 GeV/n up to 10^5 GeV/n with single-element resolution. Spectra for elements from oxygen to iron as measured by TRACER during an Antarctic circumpolar flight in 2003 are compiled in Fig. 1²¹. The data extend previous results from the space experiments HEAO (satellite) and CRN (on the Space Shuttle) to higher energies. Currently, the data of a flight from Sweden to Alaska in 2006 are analyzed. TRACER has been flown with an upgraded electronics being able to record elements from boron to iron. The new data are expected to extend measurements of the boron-to-carbon ratio up to 10^5 GeV/n, thus, providing, unprecedented information of the propagation properties approaching the knee in the energy spectrum.

3 Indirect measurements

Air shower experiments extend the findings of direct measurements to highest energies. They provide information on the all-particle energy spectrum, the (average) mass composition, and, recently, information on energy spectra for elemental groups. These results contribute to the understanding of the origin of structures in the all-particle spectrum, like the knee or the second knee.

When high-energy cosmic-ray particles penetrate the Earth's atmosphere they interact and generate a cascade of secondary particles, the extensive air showers. Two types of experiments may be distinguished to register air showers: installations measuring the longitudinal development of showers (or the depth of the shower maximum) in the atmosphere by observations of




Figure 1: Energy spectra for individual elements as measured by the TRACER, HEAO, and CRN experiments²¹.




Figure 2: Energy spectra for primary protons and helium nuclei, unfolded from air showers registered by KASCADE within three zenith angle intervals²².

Čerenkov or fluorescence light and apparatus measuring the density (and energy) of secondary particles (electrons, muons, hadrons) at ground level.

The arrival direction of cosmic rays is determined by measurements of the arrival times of secondary particles at the detectors. The shower energy is proportional to the total light collected or to the total number of particles recorded at observation level. More challenging is an estimation of the mass of the primary particle. It is either derived by a measurement of the depth of the shower maximum X_{max} or measuring the electron-to-muon ratio (N_e/N_μ) in showers. A Heitler model of hadronic showers^{23,24,11} can be used to show that air shower experiments are only sensitive to the logarithm of the mass of the primary particle ($\ln A$). To measure the composition with a resolution of 1 unit in $\ln A$ the shower maximum has to be measured to an accuracy of about 37 g/cm^2 or the N_e/N_μ ratio has to be determined with a relative error around 16%. Due to the large intrinsic fluctuations in air showers, with existing experiments at most groups of elements can be reconstructed with $\Delta \ln A \approx 0.8 - 1$.

The all-particle energy spectrum up to about 10^{18} eV is reasonably well known. The spectra obtained by different experiments agree well in shape and absolute flux taking a $\pm 10\%$ uncertainty in the energy determination of the individual experiments into account⁶. The spectrum can be described by a power law $dN/dE \propto E^\gamma$, with an spectral index changing at the knee ($E_k = 4.5 \text{ PeV}$) from $\gamma = -2.7$ to $\gamma = -3.1$.

For the mean (logarithmic) mass of cosmic rays there seems to be a dependence on the technique applied for the air shower observations²⁵. Two classes of experiments may be distinguished: detectors measuring the depth of the shower maximum and experiments measuring (the lateral distribution of) particles at ground level. This indicates inconsistencies in the interaction models used to interpret the data. For example, introducing lower inelastic cross sections in the interaction model QGSJET 01 and slightly increasing the elasticity, good agreement between the two classes of experiments can be achieved²⁵. Interpreting the data with this model yields a consistent picture with a mean logarithmic mass smoothly continued from direct measurements up to about 10^{15} eV , then exhibiting an increase up to energies around 10^{17} eV .

A big step forward in understanding the origin of the knee in the all-particle energy spectrum are the results of the KASCADE experiment. The latter comprises a $200 \times 200 \text{ m}^2$ scintillator array to register the electromagnetic and muonic component²⁶, a hadron calorimeter equipped

with liquid ionization chambers²⁷, and a limited-streamer tube muon tracking detector²⁸. The simultaneous detection of different shower components allows detailed studies of hadronic interaction models^{29,30,31,32,33,34,35}. The most consistent models are then applied to unfold energy spectra for five groups of elements from the measured air shower data³⁶. The energy spectra of the light elements (protons and helium) exhibit a depression of the flux at high energies while the spectra of the heavier groups continue to follow power laws to higher energies. The results obtained using different hadronic interaction models to interpret the data agree qualitatively. However, there are differences in the absolute flux obtained, indicating inconsistencies in the interaction models. These results have recently been confirmed by an analysis taking also inclined showers into account²². As example, the spectra obtained for primary protons and helium nuclei for three different intervals of the cosmic-ray arrival direction are shown in Fig. 2. The different data sets are consistent with each other.

Similar energy spectra for groups of elements have been obtained by the EAS-TOP³⁷, GRAPES-3³⁸, and Tibet³⁹ experiments¹⁰. Overall, a consistent 'standard picture' evolves^{6,10}: the energy spectra as obtained by direct measurements can be extrapolated to high energies, assuming power laws. The data are compatible with cut-offs in the spectra proportional to their nuclear charge Z . The all-particle spectrum and mean logarithmic mass thus obtained are compatible with air shower data. The combined energy spectra for groups of elements obtained from direct and indirect measurements extend over seven orders of magnitude in energy. They are compatible with contemporary models of the acceleration of cosmic rays in supernova remnants and models of the diffusive propagation of the particles in the Galaxy¹⁰.

Another important aspect is the anisotropy of the arrival direction of cosmic rays, giving hints to understand the propagation of the particles in the Galaxy. If the knee is caused by leakage of particles from the Galaxy, one would expect less particles arriving from directions perpendicular to the galactic disk. Frequently, the Rayleigh formalism is applied to quantify the observations⁴⁰. Upper limits as obtained by KASCADE and preliminary results obtained by the KASCADE-Grande experiment⁴¹ indicate an increase of the anisotropy amplitudes as function of energy. However, an interpretation has to take into account that the available statistics decreases roughly as $\propto E^{-3}$, therefore, an increase of the observed anisotropy is expected, which is of the order of the observed increase. The observed anisotropy amplitude is compatible with a diffusion model of the particle propagation^{40,42,43}.

4 TeV γ -ray astronomy

Observation of TeV γ -rays with atmospheric Čerenkov telescopes, like the H.E.S.S.⁴⁵, MAGIC⁴⁶, or VERITAS⁴⁷ experiments, give the most direct hints towards the acceleration sites of cosmic rays.

The energy density of galactic cosmic rays amounts to about $\rho_E \approx 1 \text{ eV/cm}^3$. The power required to sustain a constant cosmic-ray intensity can be estimated as $L_{cr} = \rho_{cr} V / \tau_{esc} \approx 10^{41} \text{ erg/s}$, where τ_{esc} is the residence time of cosmic rays in a volume V (the Galaxy, or the galactic halo). With a rate of about three supernovae per century in a typical Galaxy the energy required could be provided by a small fraction ($\approx 10\%$) of the kinetic energy released in supernova explosions. This has been realized already by Baade and Zwicky⁴⁸. The actual mechanism of acceleration remained mysterious until Fermi⁴⁹ proposed a process that involved interaction of particles with large-scale magnetic fields in the Galaxy. Eventually, this led to the currently accepted model of cosmic-ray acceleration by a first-order Fermi mechanism that operates in strong shock fronts which are powered by the explosions and propagate from the supernova remnant (SNR) into the interstellar medium^{50,51,52,53}.

The theory of acceleration of (hadronic) cosmic rays at shock fronts in supernova remnants is strongly supported by recent measurements of the H.E.S.S. experiment^{54,44}, observing TeV




Figure 3: The SNR RX J1713.7-3946 as seen by the H.E.S.S. experiment, spatially resolved in TeV γ -ray light. *Left*: γ -ray excess contours, superimposed are 14 fields for which spectra have been obtained independently. The corresponding integral flux is displayed as function of photon spectral index (*right*). For details see ⁴⁴.

γ -rays from the shell type supernova remnant RX J1713.7-3946, originating from a core collapse supernova of type II/Ib. For the first time, a SNR could be spatially resolved in γ -rays and spectra have been derived directly at a potential cosmic-ray source as can be seen in Fig. 3. The figure shows on the left-hand side excess contours of TeV γ -rays. The SNR has been divided into 14 fields for which spectra have been obtained independently⁴⁴. The corresponding integral flux values are plotted against the spectral index on the right-hand panel. The measurements yield a spectral index $\gamma = -2.19 \pm 0.09 \pm 0.15$ for the observed γ -ray flux averaged over the complete source.

A model for the acceleration of hadronic particles in the SNR is supported by measurements in various energy ranges from radio wavelengths to TeV γ -rays⁵⁵. An important feature of the model is that efficient production of nuclear cosmic rays leads to strong modifications of the shock with large downstream magnetic fields ($B \approx 100 \mu\text{G}$). Due to this field amplification the electrons are accelerated to lower maximum energy and for the same radio/x-ray flux less electrons are needed. Consequently, the inverse Compton and Bremsstrahlung fluxes are relatively low only. The model predicts that the spectrum of decaying neutral pions, generated in interactions of accelerated hadrons with material in the vicinity of the source, is clearly dominant over electromagnetic emission generated by inverse Compton effect and non-thermal Bremsstrahlung. The measurements are compatible with a nonlinear kinetic theory of cosmic-ray acceleration in supernova remnants and imply that this supernova remnant is an effective source of nuclear cosmic rays, where about 10% of the mechanical explosion energy are converted into nuclear cosmic rays^{55,56}.

Further quantitative evidence for the acceleration of hadrons in supernova remnants is provided by measurements of the HEGRA experiment⁵⁷ of TeV γ -rays from the SNR Cassiopeia A⁵⁸ and by measurements of the H.E.S.S. experiment from the SNR "Vela Junior" (SNR RX J0852.0-4622)⁵⁹.

For completeness, it should also be mentioned that the detection of γ -rays with GeV energies contributed to the understanding of cosmic-ray propagation. The diffuse γ -ray background detected with the EGRET satellite experiment⁶⁰ exhibits a structure in the GeV region, which is interpreted as indication for the interaction of propagating cosmic rays with interstellar matter⁶¹. This is the most direct evidence for cosmic rays propagating in the galactic halo, well outside the galactic disc.

5 Conclusions and outlook

In the last decade our understanding of the origin of galactic cosmic rays has been significantly improved by multidisciplinary efforts, combining key observations of the direct and indirect measurements of cosmic rays as well as the detection of γ -rays. The observations by the H.E.S.S. γ -ray telescope give clear hints that hadronic particles are accelerated in SNR. The data are compatible with a model of first order Fermi acceleration at strong shock fronts. The particles propagate in a diffusive process through the Galaxy. Parameters of the propagation models have been constrained by direct measurements above the atmosphere. The KASCADE experiment has shown that the energy spectra of the light elements exhibit a cut-off structure, while the spectra of heavier elemental groups follow power laws to higher energies. The observed spectra seem to be compatible with the assumption of power laws and a cut-off energy proportional to the nuclear charge. This implies that the knee in the all-particle energy spectrum is caused by a cut-off of the light elements. The shape of the all-particle spectrum at higher energies is then determined by the subsequent cut-offs of all elemental species in cosmic rays. Most likely, the astrophysical origin of the knee is a combination of the maximum energy reached in the acceleration process and leakage from the Galaxy during propagation⁶². More exotic ideas about the cause of the knee are most likely excluded^{9,10}.

In conclusion this gives a qualitative 'standard picture' of the origin of galactic cosmic rays. However, several details remain unclear and a precise quantitative description of all aspects of the acceleration and propagation mechanisms is still missing. Among the open questions are: It is not clear how to precisely match the spectral indices observed at Earth to the spectra at the sources, being compatible with the TeV γ -ray observations and the modification of the spectral slope during propagation.

A precise astrophysical interpretation of air shower data is presently limited by the understanding of hadronic interactions in the atmosphere, thus the exact shape of the energy spectra at their corresponding knees is unknown.

Contemporary assumptions on the parameters of cosmic-ray propagation models yield anisotropies in the arrival directions not observed by air shower experiments^{63,62}.

In the near future experiments like TRACER aim to reveal details of the cosmic-ray propagation for energies approaching the knee⁶⁴. Experiments at the LHC probing the extreme forward direction of phase space will improve the description of high-energy hadronic interactions⁶⁵. Also the exploration of the end of the galactic component and the transition to extragalactic particles in the energy range from $10^{17} - 10^{18}$ eV will be of importance. Key experiments in this energy region are KASCADE-Grande (a 0.5 km² extension of the KASCADE experiment)⁶⁶, Ice Cube/ Ice Top (a 1 km² air shower experiment and neutrino telescope at the South Pole)^{67,68}, and HEAT/AMIGA (a 25 km² extension of the Pierre Auger Observatory to lower energies)⁶⁹.

Around 10^{18} eV two features appear in the all-particle energy spectrum. The second knee at $E_{2nd} \approx 400 \text{ PeV} \approx 92 \times E_k$, where the spectrum exhibits a steepening to $\gamma \approx -3.3$, and the ankle at about 4 EeV, above this energy the spectrum seems to flatten again to $\gamma \approx -2.7$. The region around 4 EeV is sometimes also called "the dip" in the spectrum.

A possible cause for the second knee is the end of the galactic component, when all elements successively have reached their cut-off energies, the latter being proportional to their nuclear charge^{6,62}. If one assumes that ultra-heavy elements (heavier than iron) play an important role to understand the second knee, the factor of 92 between the energies of the knee and the second knee can be easily understood as the nuclear charge of the heaviest elements in the periodic table.

The dip is proposed to be caused by electron-positron pair production of cosmic rays on photons of the cosmic microwave background⁷⁰.

The investigation of the transition region and a precise measurement of the galactic all-

particle spectrum is also important for an estimate of the energy content of extragalactic cosmic rays. For example, the extragalactic component needed according to the poly-gonato model to sustain the observed all-particle flux at highest energies has an energy density of $\rho_E^{exg} = 3.7 \cdot 10^{-7} \text{ eV/cm}^3$.

A promising rediscovered technique for the exploration of cosmic rays with energies above 10^{16} eV is the detection of radio signals from air showers⁷¹. Most likely, the emission mechanism is coherent synchrotron radiation of electrons with energies slightly lower than the critical energy ($E_{krit} = 85 \text{ MeV}$ in air) deflected in the magnetic field of the Earth (geosynchrotron radiation)⁷². The LOPES experiment, registering showers in coincidence with the KASCADE-Grande experiment has demonstrated the feasibility of this approach⁷³. Radio detection of air showers is also pursued in the CODALEMA experiment⁷⁴ and with the LOFAR radio telescope⁷⁵. First radio pulses from air showers have been recorded as well with antennae set up at the southern site of the Pierre Auger Observatory⁷⁶.

Acknowledgments

I would like to thank the organizers of the XXth Rencontres de Blois for their invitation to this very inspiring scientific meeting. I'm grateful to my colleagues from TRACER, KASCADE-Grande, LOPES, LOFAR, and the Pierre Auger Observatory for fruitful discussions.

Reference

1. V. Hess, *Phys. Zeitschr.* **13**, 1084 (1912).
2. M. Nagano *et al.*, *Rev. Mod. Phys.* **72**, 689 (2000).
3. J. Hörandel, *Rev. Mod. Astron.* **20**, 203 (2008).
4. F. Halzen, astro-ph/0604441 (2006).
5. T. Gaisser, *Cosmic Rays and Particle Physics*, Cambridge University Press (1999).
6. J. Hörandel, *Astropart. Phys.* **19**, 193 (2003).
7. J. Hörandel, *Astropart. Phys.* **21**, 241 (2004).
8. J. Hörandel, *Il Nuovo Cimento B* **120**, 825 (2005).
9. J. Hörandel, *Int. J. Mod. Phys. A* **20**, 6753 (2005).
10. J. Hörandel, *Adv. Space Res.* **41**, 442 (2008).
11. J. Hörandel, *Nucl. Instr. & Meth. A* **588**, 181 (2008).
12. M. Wiedenbeck *et al.*, *Proc. 28th Int. Cosmic Ray Conf., Tsukuba* **4**, 1899 (2003).
13. N. Yanasak *et al.*, *Astrophys. J.* **563**, 768 (2001).
14. M. Garcia-Munoz *et al.*, *Astrophys. J.* **201**, L141 (1975).
15. J. Simpson *et al.*, *Space Sci. Rev.* **46**, 205 (1988).
16. M. Garzia-Munoz *et al.*, *Astrophys. J. Suppl. Ser.* **64**, 269 (1987).
17. W. Jones, *Proc. 29th Int. Cosmic Ray Conf., Pune* **10**, 173 (2005).
18. T. Guzik *et al.*, *Adv. Space Res.* **33**, 1763 (2004).
19. E. Seo *et al.*, *Adv. Space Res.* **33**, 1777 (2004).
20. F. Gahbauer *et al.*, *Astrophys. J.* **607**, 333 (2004).
21. P. Boyle *et al.*, *Proc. 30th Int. Cosmic Ray Conf., Merida* in press (2007); M. Ave *et al.*, *Astrophys. J.* **678**, 262 (2008).
22. H. Ulrich *et al.*, *Proc. 30th Int. Cosmic Ray Conf., Merida* in press (2007).
23. J. Matthews, *Astropart. Phys.* **22**, 387 (2005).
24. J. Hörandel, *Mod. Phys. Lett. A* **22**, 1533 (2007).
25. J. Hörandel, *J. Phys. G: Nucl. Part. Phys.* **29**, 2439 (2003).
26. T. Antoni *et al.*, *Nucl. Instr. & Meth. A* **513**, 490 (2003).
27. J. Engler *et al.*, *Nucl. Instr. & Meth. A* **427**, 528 (1999).

28. P. Doll *et al.*, *Nucl. Instr. & Meth. A* **488**, 517 (2002).
29. T. Antoni *et al.*, *J. Phys. G: Nucl. Part. Phys.* **25**, 2161 (1999).
30. T. Antoni *et al.*, *J. Phys. G: Nucl. Part. Phys.* **27**, 1785 (2001).
31. J. Milke *et al.*, *Acta Physica Polonica B* **35**, 341 (2004).
32. J. Milke *et al.*, *Proc. 29th Int. Cosmic Ray Conf., Pune* **6**, 125 (2005).
33. W. Apel *et al.*, *J. Phys. G: Nucl. Part. Phys.* **34** (2007) 2581.
34. J. Hörandel *et al.*, *Proc. 29th Int. Cosmic Ray Conf., Pune* **6**, 121 (2005).
35. J. Hörandel, *Nucl. Phys. B (Proc. Suppl.)* **122**, 455 (2003).
36. T. Antoni *et al.*, *Astropart. Phys.* **24**, 1 (2005).
37. G. Navarra *et al.*, *Proc. 28th Int. Cosmic Ray Conf., Tsukuba* **1**, 147 (2003).
38. Y. Hayashi *et al.*, *Proc. 29th Int. Cosmic Ray Conf., Pune* **10**, 243 (2005).
39. M. Amenomori *et al.*, *Phys. Rev. D* **62**, 112002 (2000).
40. T. Antoni *et al.*, *Astrophys. J.* **604**, 687 (2004).
41. S. Over *et al.*, *Proc. 30th Int. Cosmic Ray Conf., Merida* in press (2007).
42. G. Maier *et al.*, *Int. J. Mod. Phys. A* **20**, 6840 (2005).
43. J. Candia *et al.*, *J. Cosmol. Astropart. Phys.* **5**, 3 (2003).
44. F. Aharonian *et al.*, *Astron. & Astroph.* **449**, 223 (2006).
45. J. Hinton, *New Astronomy Reviews* **48**, 331 (2004).
46. D. Ferenc *et al.*, *Nucl. Instr. & Meth. A* **553**, 274 (2005).
47. T. Weekes *et al.*, *Astropart. Phys.* **17**, 221 (2002).
48. W. Baade *et al.*, *Phys. Rev.* **46**, 76 (1934).
49. E. Fermi, *Phys. Rev.* **75**, 1169 (1949).
50. W. Axford *et al.*, *Proc. 15th Int. Cosmic Ray Conf., Ploudiv* **11**, 132 (1977).
51. G. Krymsky, *Dok. Acad. Nauk USSR* **234**, 1306 (1977).
52. A. Bell, *Mon. Not. R. Astron. Soc.* **182**, 147 (1978).
53. R. Blanford *et al.*, *Astrophys. J.* **221**, L29 (1978).
54. F. Aharonian *et al.*, *Nature* **432**, 75 (2004).
55. H. Völk *et al.*, *Astron. & Astroph.* **451**, 981 (2006).
56. E. Berezhko *et al.*, *Astrophys. J.* **661**, L175 (2007).
57. F. Aharonian *et al.*, *Astron. & Astroph.* **370**, 112 (2001).
58. E. Berezhko *et al.*, *Astron. & Astroph.* **400**, 971 (2003).
59. H. Völk, astro-ph/0603502 (2006).
60. A. Strong *et al.*, *Astron. & Astroph.* **308**, L21 (1996).
61. A. Strong *et al.*, preprint astro-ph/9903370 (1999).
62. J. Hörandel *et al.*, *Astropart. Phys.* **27**, 119 (2007).
63. V. Ptuskin, *Adv. Space Res.* **19**, 697 (1997).
64. D. Müller *et al.*, *Proc. 30th Int. Cosmic Ray Conf., Merida* in press (2007).
65. R. Engel, *Nucl. Phys. B (Proc. Suppl.)* **151**, 437 (2006).
66. G. Navarra *et al.*, *Nucl. Instr. & Meth. A* **518**, 207 (2004).
67. M. Kestel *et al.*, *Nucl. Instr. & Meth. A* **535**, 139 (2004).
68. T. Gaisser *et al.*, *Proc. 28th Int. Cosmic Ray Conf., Tsukuba* **2**, 1117 (2003).
69. H. Kalges *et al.*, *Proc. 30th Int. Cosmic Ray Conf., Merida* in press (2007).
70. V. Berezhinsky, astro-ph/0509069 (2005).
71. H. Allan, *Progress in Elementary Particles and Cosmic Ray Physics*, pg. 169, J.G. Wilson & S.G. Wouthuysen eds., North Holland (1971).
72. T. Huege *et al.*, *Astropart. Phys.* **24**, 116 (2005).
73. H. Falcke *et al.*, *Nature* **435**, 313 (2005).
74. D. Ardouin *et al.*, *Nucl. Instr. & Meth. A* **555**, 148 (2005).
75. LOFAR, www.lofar.org (2007).
76. A. van den Berg *et al.*, *Proc. 30th Int. Cosmic Ray Conf., Merida* in press (2007).